

AD-A128 799

ESTIMATION FOR DIFFUSION PROCESSES UNDER MISSPECIFIED  
MODELS(U) FLORIDA STATE UNIV TALLAHASSEE DEPT OF  
STATISTICS I W MCKEAGUE MAR 83 FSU-STATISTICS-M653  
ARO-19367.4-MA DAAG29-82-K-0168


1/1

UNCLASSIFIED

F/G 12/1

NL


END  
DATE  
INDEXED  
NO. OF  
DTIC



MICROCOPY RESOLUTION TEST CHART  
NATIONAL BUREAU OF STANDARDS-1963-A

ARO 19367.4-MA

(12)

AD A 128 799

**ESTIMATION FOR DIFFUSION PROCESSES  
UNDER MISSPECIFIED MODELS**


by

**Ian W. McKeague**

**FSU Statistics Report No. M653  
USARO Technical Report No. D-59**

**March, 1983**

**The Florida State University  
Department of Statistics  
Tallahassee, Florida 32306**



**Research supported by the U.S. Army Research Office Grant DAAG 29-82-K-0168.**

**Key Words: Diffusion processes, maximum likelihood estimates, misspecified models.**

This document has been approved  
for public release and sale; its  
distribution is unlimited.

DTIC FILE COPY

83 06 01 01

**ESTIMATION FOR DIFFUSION PROCESSES  
UNDER MISSPECIFIED MODELS**

by

**Ian W. McKeague  
Florida State University**

**ABSTRACT**

The asymptotic behavior of the maximum likelihood estimator of a parameter in the drift term of a stationary ergodic diffusion process is studied under conditions in which the true drift function and true noise function do not coincide with those specified by the parametric model.



SEARCHED	<input checked="" type="checkbox"/>
SERIALIZED	<input type="checkbox"/>
INDEXED	<input type="checkbox"/>
FILED	<input type="checkbox"/>

1983 JUN 01 01

A

83 06 01 01

## 1. Introduction.

Consider the problem of estimating the drift function  $b(x)$  of a stationary diffusion process  $(X_t)$  given by

$$dX_t = b(X_t)dt + \sigma(X_t)dW_t, \quad t \geq 0,$$

where the process is observed over  $[0, T]$ . The method of maximum likelihood can be used if  $b(x)$  is assumed to have a parametric form  $f(x, \theta)$ ,  $\theta \in \Theta$ . Brown and Hewitt (1975), Kutoyants (1977), Lanská (1979) and Prakasa Rao and Rubin (1981) have shown that the maximum likelihood estimator of  $\theta$  is consistent and asymptotically normal. Nonparametric methods of estimating  $b$  have been developed by Banon (1978) and Geman (1980).

Suppose that a parametric model for the process  $(X_t)$  is given by

$$dX_t = f(X_t, \theta)dt + \gamma(X_t)dW_t, \quad t \geq 0.$$

This paper studies the asymptotic behavior of the maximum likelihood estimator of  $\theta$  under departures of the true drift function  $b(x)$  or true noise function  $\sigma(x)$  from those specified by the parametric model.

The need for such analysis stems from the desirability of using estimators that are robust under small departures from the underlying model. This kind of analysis is familiar in other settings, for example Huber (1967), White (1981) and Berger and Langberg (1981).

In Section 2 it is shown that the maximum likelihood estimator converges almost surely to a parameter  $\theta^*$  such that  $f(x, \theta^*)$  minimizes an  $L^2$  distance of the parametric family from the true drift function. The  $L^2$  distance is defined with respect to the measure  $\gamma^{-2}(x)d\nu(x)$ , where  $\nu$  is the stationary distribution of the process. Asymptotic normality is also established. In Section 3 we discuss a way of estimating the difference in goodness of fit of two separate parametric families to the true drift function. Examples are given at the end of each section.

## 2. Maximum likelihood estimation under misspecified models.

Let  $(X_t, t \geq 0)$  be a stationary, ergodic process which is assumed to be the unique solution of the stochastic differential equation

$$(2.1) \quad dX_t = b(X_t)dt + \sigma(X_t)dW_t, \quad t \geq 0$$

where  $X_0$  is distributed according to the stationary distribution of the process,  $b$  and  $\sigma$  are unknown measurable functions and  $(W_t, t \geq 0)$  is a standard Wiener process. Assume that  $(X_t)$  has inaccessible boundaries on the state space  $(-\infty, \infty)$ . Using the notation of Mandl (1968), let

$$B(x) = 2 \int_0^x \frac{b(y)}{\sigma^2(y)} dy, \quad p(x) = \int_0^x \exp(-B(y)) dy,$$

$$m(x) = 2 \int_0^x \frac{\exp B(y)}{\sigma^2(y)} dy,$$

where the integrals are assumed to exist. Provided that  $m(+\infty) < \infty$  and  $m(-\infty) > -\infty$ , the stationary distribution, denoted  $\nu$ , has distribution function  $M^{-1}m(x)$  where  $M = m(+\infty) - m(-\infty)$ .

Suppose a parametric model is used to estimate the drift function  $b(x)$  by the method of maximum likelihood. Let  $\Theta$  denote a closed bounded interval. A family of measurable drift functions  $\{f(x, \theta), \theta \in \Theta\}$  and a measurable noise function  $\gamma(x) > 0$  are provided and inference is based on the model

$$(2.2) \quad dX_t = f(X_t, \theta)dt + \gamma(X_t)dW_t, \quad t \geq 0.$$

The process  $(X_t)$  is observed over  $[0, T]$ . Let  $\mu_\theta^T$  and  $\mu^T$  denote the measures induced on  $C[0, T]$  by process satisfying (2.2) and the process

$$dY_t = \gamma(Y_t)dW_t, \quad t \geq 0$$

$$Y_0 = X_0,$$

respectively. Under conditions given in Liptser and Shirayev (1977, Theorem 7.19) it follows that  $\mu_\theta^T \ll \mu^T$  for all  $\theta \in \Theta$  and the log likelihood function  $l_T(\theta) = \log[du_\theta^T/du^T](X)$  is given by

$$(2.3) \quad l_T(\theta) = \int_0^T \frac{f(X_t, \theta)}{\gamma^2(X_t)} dX_t - \frac{1}{2} \int_0^T \left[ \frac{\bar{f}(X_t, \theta)}{\gamma(X_t)} \right]^2 dt, \quad \text{a.e. } (\mu^T).$$

A maximum likelihood estimator calculated from  $l_T(\theta)$  is denoted  $\hat{\theta}_T$ .

Assume that  $E \left[ \frac{\bar{b}(X_0) - f(X_0, \theta)}{\gamma(X_0)} \right]^2 < \infty$  for all  $\theta \in \Theta$  and, as a function of

$\theta$ , has a unique minimum at  $\theta^* \in \Theta$ . The following results describe the asymptotic behavior of  $\hat{\theta}_T$  when the observed process satisfies (2.1). The conditions are stated later.  $g'$ ,  $g''$  denote first and second partial derivatives of a function  $g(x, \theta)$  with respect to  $\theta$ .

Theorem 2.1. Under conditions (C1)-(C3),  $\hat{\theta}_T \rightarrow \theta^*$  a.s. as  $T \rightarrow \infty$ .

Theorem 2.2. Under conditions (C1)-(C7),  $T^{1/2}(\hat{\theta}_T - \theta^*) \xrightarrow{D} N(0, \Sigma)$  where

$$(2.4) \quad \Sigma = \frac{2M \int_{-\infty}^{\infty} g'(y, \theta^*) \int_y^0 \int_{-\infty}^s g'(z, \theta^*) dm(z) dp(s) dm(y)}{\left\{ \int_{-\infty}^{\infty} g''(x, \theta^*) dm(x) \right\}^2},$$

$$(2.5) \quad g(x, \theta) = \left[ \frac{\bar{f}(x, \theta)}{\gamma(x)} \right]^2 + \sigma^2(x) \frac{\partial}{\partial x} \left( \frac{f(x, \theta)}{\gamma^2(x)} \right).$$

Theorem 2.3. In the special case that the drift function has been correctly specified, i.e.  $b(x) = f(x, \theta_0)$  for some  $\theta_0 \in \Theta$ , then (whether or not the noise function has been correctly specified)

(i)  $\hat{\theta}_T \rightarrow \theta_0$  a.s. as  $T \rightarrow \infty$  under conditions (C1)-(C3);

(11)  $T^h(\hat{\theta}_T - \theta_0) \xrightarrow{D} N(0, V)$  under conditions (C1)-(C9), where

$$(2.6) \quad V = \frac{\int_{-\infty}^{\infty} \left[ \frac{f'(y, \theta_0)}{\gamma(y)} \right]^2 \left[ \frac{\sigma(y)}{\gamma(y)} \right]^2 d\nu(y)}{\left\{ \int_{-\infty}^{\infty} \left[ \frac{f'(x, \theta_0)}{\gamma(x)} \right]^2 d\nu(x) \right\}^2}.$$

Remarks.

(a) The parameter  $\theta^*$  can often be determined from the moments of the stationary distribution. In general it is difficult to evaluate  $\Sigma$  unless the drift function has been correctly specified (as in Theorem 2.3). Some examples in which  $\theta^*$  and  $\Sigma$  are explicitly calculated are given at the end of this section.

(b) When both drift and noise functions have been correctly specified  $V$  reduces to the previously known formula

$$V_0 = \left( \int_{-\infty}^{\infty} \left[ \frac{f'(x, \theta_0)}{\sigma(x)} \right]^2 d\mu_0(x) \right)^{-1}.$$

The Cauchy-Schwarz inequality yields  $V \geq V_0$ .

(c) Theorem 2.3 (i) can be interpreted as a robustness result for the maximum likelihood estimator;  $\hat{\theta}_T$  remains consistent under misspecifications of the noise function.

(d) Laska (1979) introduced a minimum contrast estimator  $\tilde{\theta}_T$  which minimizes

$$\int_0^T h(X_t, \theta) dt$$

where



$$h(x, \theta) = \left[ \frac{f(x, \theta)}{\gamma(x)} \right]^2 + \gamma^2(x) \frac{\partial}{\partial x} \left( \frac{f(x, \theta)}{\gamma^2(x)} \right).$$

If  $h$  satisfies the conditions (C1)-(C7) in place of  $g$  and the function  $Eh(X_0, \theta)$  has a unique minimum at  $\theta_1$ , then using similar proofs to Theorems 2.1 and 2.2,  $\tilde{\theta}_T \rightarrow \theta_1$  a.s. and  $T^{1/2}(\tilde{\theta}_T - \theta_1) \xrightarrow{D} N(0, \Sigma_1)$ , where  $\Sigma_1$  is given by (2.4) with  $g$  replaced by  $h$ . Even when the drift function is correctly specified, it is possible for  $\theta_1 \neq \theta_0$  when the noise function is misspecified, so that  $\tilde{\theta}_T$  can fail to be a consistent estimator of  $\theta_0$  while  $\hat{\theta}_T$  remains consistent.

Conditions.

$$(C1) \quad |f(x, \theta_1) - f(x, \theta_2)| \leq J(x)\psi(\theta_1 - \theta_2), \quad x \in R, \quad \theta_1, \theta_2 \in \Theta, \text{ where} \\ E \left[ \frac{J(X_0)}{\gamma(X_0)} \right]^2 < \infty \text{ and } \lim_{\alpha \rightarrow 0} \psi(\alpha) = 0.$$

$$(C2) \quad E \left[ \frac{\sigma(X_0)}{\gamma(X_0)} \right]^2 < \infty \text{ and } |f(x, \theta)| \leq K(x), \quad x \in R, \quad \theta \in \Theta, \text{ where} \\ E \left[ \frac{K(X_0)}{\gamma(X_0)} \right]^2 < \infty.$$

$$(C3) \quad f(x, \theta) \text{ is continuous in } (x, \theta) \text{ and differentiable with respect} \\ \text{to } \theta. \text{ There exists } \alpha > 0 \text{ such that } |f'(x, \theta_1) - f'(x, \theta_2)| \leq c(x)|\theta_1 - \theta_2|^\alpha, \\ x \in R, \quad \theta_1, \theta_2 \in \Theta, \text{ where } E \left[ \frac{\sigma(X_0)c(X_0)}{\gamma^2(X_0)} \right]^2 < \infty.$$

$$(C4) \quad f(x, \theta)\gamma^{-2}(x) \text{ has a continuous first partial derivative with} \\ \text{respect to } x \text{ for each } \theta \in \Theta.$$

$$(C5) \quad \lim_{x \rightarrow \pm\infty} f(x, \theta)\gamma^{-2}(x)\exp B(x) = 0, \quad \forall \quad \theta \in \Theta.$$

(C6) The partial derivatives  $g'$ ,  $g''$ ,  $G'$ ,  $G''$  exist and are continuous in  $(x, \theta)$ , where  $G(x, \theta)$  is defined by

$$(2.7) \quad G(x, \theta) = \int_0^x \frac{f(y, \theta)}{\gamma^2(y)} dy.$$

(C7)  $|g''(x, \theta_1) - g''(x, \theta_2)| \leq U(x)\phi(\theta_1 - \theta_2)$ ,  $x \in R$ ,  $\theta_1, \theta_2 \in \Theta$ , where  $E U(X_0) < \infty$  and  $\lim_{\alpha \rightarrow 0} \phi(\alpha) = 0$ .

$$(C8) \quad \lim_{x \rightarrow \pm\infty} f'(x, \theta_0) \gamma^{-2}(x) [\exp B(x)] \int_0^x f'(s, \theta_0) \gamma^{-2}(s) ds = 0.$$

$$(C9) \quad \lim_{x \rightarrow \pm\infty} f''(x, \theta_0) \gamma^{-2}(x) \exp B(x) = 0.$$

(C10) Condition (C7) with  $g'$  in place of  $g''$ .

Proof of Theorem 2.1. The proof of strong consistency of  $\hat{\theta}_T$  in the correctly specified case given by Prakasa Rao and Rubin (1981, Theorem 4.1) needs only minor modifications to show that  $\hat{\theta}_T \rightarrow \theta^*$  a.s. as  $T \rightarrow \infty$  in the misspecified case. The details are given here for completeness. From (2.1) and (2.3)  $L_T(\theta)$  can be written

$$(2.6) \quad L_T(\theta) = -\frac{1}{2} \int_0^T \left[ \frac{f(X_t, \theta) - b(X_t)}{\gamma(X_t)} \right]^2 dt + \frac{1}{2} \int_0^T \left[ \frac{b(X_t)}{\gamma(X_t)} \right]^2 dt + \int_0^T \frac{f(X_t, \theta) \sigma(X_t)}{\gamma^2(X_t)} dW_t.$$

$$\text{Denote } I_T(\theta) = \int_0^T \left[ \frac{f(X_t, \theta) - b(X_t)}{\gamma(X_t)} \right]^2 dt. \text{ For } \theta_1, \theta_2 \in \Theta,$$

$$\begin{aligned} |I_T(\theta_1) - I_T(\theta_2)| &\leq \int_0^T \frac{|f(X_t, \theta_1) - f(X_t, \theta_2)| (|f(X_t, \theta_1)| + |f(X_t, \theta_2)|)}{\gamma^2(X_t)} dt \\ &\quad + 2 \int_0^T \frac{|b(X_t)| |f(X_t, \theta_1) - f(X_t, \theta_2)|}{\gamma^2(X_t)} dt \\ &\leq 2\phi(\theta_1 - \theta_2) \int_0^T \frac{J(X_t)(K(X_t) + b(X_t))}{\gamma^2(X_t)} dt, \end{aligned}$$

using conditions (C1) and (C2). By the ergodic theorem

$$\frac{1}{T} \int_0^T \frac{J(X_t)(K(X_t)+b(X_t))}{\gamma^2(X_t)} dt \rightarrow E \left[ \frac{J(X_0)(K(X_0)+b(X_0))}{\gamma^2(X_0)} \right] \text{ a.s.}$$

as  $T \rightarrow \infty$ , using conditions (C1) and (C2) again. Thus, there exists a r.v.  $C^*$ , which is finite a.s. and does not depend on  $\theta$ , such that

$$|I_T(\theta_1) - I_T(\theta_2)| \leq C^* T \psi(\theta_1 - \theta_2), \text{ for all } \theta_1, \theta_2 \in \Theta, T \geq 0.$$

Similarly, using condition (C2) again, it follows that there exists a r.v.  $D^*$ , which is finite a.s. and does not depend on  $\theta$ , such that  $I_T(\theta) \leq D^* T$ , for all  $\theta \in \Theta$ ,  $T \geq 0$ . Thus,  $\{\frac{1}{T} I_T(\cdot), T \geq 0\}$  is equicontinuous and uniformly bounded a.s. as a family of functions of  $\theta$ . By the Arzela-Ascoli theorem this family is relatively compact (a.s.) in the space of continuous functions on  $\Theta$  provided with the supremum norm. Therefore, by the ergodic theorem

$$\frac{1}{T} I_T(\theta) \xrightarrow{\text{a.s.}} E \left[ \frac{[f(X_0, \theta) - b(X_0)]^2}{\gamma(X_0)} \right],$$

uniformly in  $\theta \in \Theta$  as  $T \rightarrow \infty$ .

Now consider the second term in (2.6). By condition (C2) and the ergodic theorem

$$\frac{1}{T} \int_0^T \left[ \frac{b(X_t)}{\gamma(X_t)} \right]^2 dt \xrightarrow{\text{a.s.}} E \left[ \frac{[b(X_0)]^2}{\gamma(X_0)} \right] \text{ as } T \rightarrow \infty.$$

Next, using Lemma 4.3 of Prakasa Rao and Rubin (1981) it follows that, under conditions (C1)-(C3),

$$\frac{1}{T} \int_0^T \frac{f(X_t, \theta) \sigma(X_t)}{\gamma^2(X_t)} dW_t \xrightarrow{\text{a.s.}} 0, \text{ uniformly in } \theta \in \Theta \text{ as } T \rightarrow \infty.$$

Thus

$$(2.7) \quad \frac{1}{T} \ell_T(\theta) \text{ a.s. } \rightarrow E \left[ \frac{f(X_0, \theta) - b(X_0)}{\gamma(X_0)} \right]^2 + E \left[ \frac{b(X_0)}{\gamma(X_0)} \right]^2,$$

uniformly in  $\theta \in \Theta$  as  $T \rightarrow \infty$ . Since the r.h.s. of (2.7) has a unique maximum at  $\theta^* \in \Theta$  and  $\hat{\theta}_T$  maximizes  $\frac{1}{T} \ell_T(\theta)$ , it is easily proved that  $\hat{\theta}_T \rightarrow \theta^*$  a.s. as  $T \rightarrow \infty$ .  $\square$

Proof of Theorem 2.2. The approach used by Prakasa Rao and Rubin (1981) to find the asymptotic distribution of  $\hat{\theta}_T$  in the correctly specified case does not extend to the misspecified case. Rather, the proof of this theorem uses the technique, introduced by Laska (1979), of expressing  $\ell_T(\theta)$  in terms of Lebesgue integrals.

The function  $G(x, \theta)$  defined in (2.7) has a continuous second partial derivative with respect to  $x$  for each  $\theta \in \Theta$  by condition (C4). Applying Itô's formula, it follows that

$$\begin{aligned} G(X_T, \theta) &= G(X_0, \theta) + \int_0^T \left[ \frac{b(X_t) f(X_t, \theta)}{\gamma^2(X_t)} + \frac{1}{2} \sigma^2(X_t) G''(X_t, \theta) \right] dt \\ &\quad + \int_0^T \frac{\sigma(X_t) f(X_t, \theta)}{\gamma^2(X_t)} dW_t. \end{aligned}$$

Then, using (2.1) and (2.3),

$$\begin{aligned} \ell_T(\theta) &= \int_0^T \frac{f(X_t, \theta) b(X_t)}{\gamma^2(X_t)} dt + \int_0^T \frac{f(X_t, \theta) \sigma(X_t)}{\gamma^2(X_t)} dW_t - \frac{1}{2} \int_0^T \left[ \frac{f(X_t, \theta)}{\gamma(X_t)} \right]^2 dt \\ (2.8) \quad &= G(X_T, \theta) - G(X_0, \theta) - \frac{1}{2} \int_0^T g(X_t, \theta) dt, \end{aligned}$$

where  $g$  is defined in (2.5). Expand  $\lambda_T^1(\theta)$  about  $\hat{\theta}_T$ ,

$$\lambda_T^1(\theta^*) = \lambda_T^1(\hat{\theta}_T) + (\theta^* - \hat{\theta}_T) \lambda_T''(\bar{\theta}_T), \text{ where } |\bar{\theta}_T - \theta^*| \leq |\hat{\theta}_T - \theta^*|.$$

Consider

$$T^{-1/2} \lambda_T^1(\theta^*) = T^{-1/2} (G'(X_T, \theta^*) - G'(X_0, \theta^*)) + T^{-1/2} \int_0^T g'(X_t, \theta^*) dt.$$

Using the stationarity of  $(X_t)$ ,

$$T^{-1/2} (G'(X_T, \theta^*) - G'(X_0, \theta^*)) \xrightarrow{P} 0 \text{ as } T \rightarrow \infty.$$

Using integration by parts and condition (C5) it can be shown that

$$(2.9) \quad E g(X_0, \theta) = E \left[ \frac{f(X_0, \theta) - b(X_0)}{\gamma(X_0)} \right]^2 - E \left[ \frac{b(X_0)}{\gamma(X_0)} \right]^2,$$

and since the right hand side of this expression is minimized at  $\theta^* \in \Theta$ , it follows that  $E g(X_0, \theta)$  is minimized at  $\theta^* \in \Theta$  and  $E g'(X_0, \theta^*) = 0$ .

Then, by Mandl (1968, p. 94)

$$T^{-1/2} \int_0^T g'(X_t, \theta^*) \xrightarrow{D} N(0, \Delta),$$

where

$$\Delta = \frac{2}{M} \int_{-\infty}^{\infty} g'(y, \theta^*) \int_y^0 \int_{-\infty}^s g'(z, \theta^*) dm(z) dp(s) dm(y).$$

Thus  $T^{-1/2} \lambda_T^1(\theta^*) \xrightarrow{D} N(0, \Delta)$ . By the ergodic theorem and condition (C6)

$$\frac{1}{T} \lambda_T''(\theta^*) \xrightarrow{P} E g''(X_0, \theta^*).$$

Next, using conditions (C6), (C7) and the fact that  $\bar{\theta}_T \rightarrow \theta^*$  a.s.

$$\frac{1}{T} (L_T''(\bar{\theta}_T) - L_T''(\theta^*)) \xrightarrow{P} 0, \text{ as } T \rightarrow \infty.$$

Thus

$$\frac{1}{T} L_T''(\bar{\theta}_T) \xrightarrow{P} E g''(X_0, \theta^*) = \frac{1}{M} \int_{-\infty}^{\infty} g''(x, \theta^*) d\mu(x),$$

as  $T \rightarrow \infty$ . We conclude that

$$T^{1/2}(\bar{\theta}_T - \theta^*) \xrightarrow{D} N(0, \Sigma),$$

where  $\Sigma$  is given in (2.4). □

Proof of Theorem 2.3. Suppose that  $b(x) = f(x, \theta_0)$ , where  $\theta_0 \in \Theta$ . Then  $\theta^* = \theta_0$  and (a) follows directly from Theorem 2.1. The proof of (b) consists in showing that  $\Sigma$  in (2.4) reduces to  $V$  given in (2.6). Note that

$$g'(x, \theta_0) = 2b(x)f'(x, \theta_0)\gamma^{-2}(x) + \sigma^2(x) \frac{\partial}{\partial x} [f'(x, \theta_0)\gamma^{-2}(x)].$$

Using integration by parts,

$$\begin{aligned} \int_{-\infty}^s \sigma^2(z) \frac{\partial}{\partial z} [f'(z, \theta_0)\gamma^{-2}(z)] d\mu(z) \\ = 2f'(s, \theta_0)\gamma^{-2}(s) \exp B(s) - 2 \int_{-\infty}^s b(z)f'(z, \theta_0)\gamma^{-2}(z) d\mu(z), \end{aligned}$$

so that

$$\int_{-\infty}^s g'(z, \theta_0) d\mu(z) = 2f'(s, \theta_0)\gamma^{-2}(s) \exp B(s),$$

and

$$\int_y^0 \int_{-\infty}^s g'(z, \theta_0) dm(z) dp(s) = 2 \int_y^0 f'(s, \theta_0) \gamma^{-2}(s) ds.$$

Using integration by parts with condition (C8),

$$\begin{aligned} & \int_{-\infty}^{\infty} \sigma^2(y) \frac{\partial}{\partial y} [f'(y, \theta_0) \gamma^{-2}(y)] \int_y^0 f'(s, \theta_0) \gamma^{-2}(s) ds dm(y) \\ &= -2 \int_{-\infty}^{\infty} f'(y, \theta_0) \gamma^{-2}(y) \frac{\partial}{\partial y} \left[ \int_y^0 f'(s, \theta_0) \gamma^{-2}(s) ds \exp B(y) \right] dy \\ &= \int_{-\infty}^{\infty} \left[ \frac{f'(y, \theta_0)}{\gamma(y)} \right]^2 \left[ \frac{\sigma(y)}{\gamma(y)} \right]^2 dm(y) - 2 \int_{-\infty}^{\infty} b(y) f'(y, \theta_0) \gamma^{-2}(y) \int_y^0 f'(s, \theta_0) \gamma^{-2}(s) ds dm(y). \end{aligned}$$

It follows that the numerator of  $\Sigma$  reduces to

$$4M \int_{-\infty}^{\infty} \left[ \frac{f'(y, \theta_0)}{\gamma(y)} \right]^2 \left[ \frac{\sigma(y)}{\gamma(y)} \right]^2 dm(y).$$

Now consider the denominator of  $\Sigma$ .

$$g''(x, \theta_0) = 2 \left[ \frac{f'(x, \theta_0)}{\gamma(x)} \right]^2 + 2f(x, \theta_0) f''(x, \theta_0) \gamma^{-2}(x) + \sigma^2(x) \frac{\partial}{\partial x} [f''(x, \theta_0) \gamma^{-2}(x)].$$

Using integration by parts and condition (C9)

$$\int_{-\infty}^{\infty} \sigma^2(x) \frac{\partial}{\partial x} [f''(x, \theta_0) \gamma^{-2}(x)] dm(x) = -2 \int_{-\infty}^{\infty} f(x, \theta_0) f''(x, \theta_0) \gamma^{-2}(x) dm(x)$$

and it follows that

$$\int_{-\infty}^{\infty} g''(x, \theta_0) dm(x) = 2 \int_{-\infty}^{\infty} \left[ \frac{f'(x, \theta_0)}{\gamma(x)} \right]^2 dm(x).$$

This completes the proof of the theorem. □

Examples.

1. This example presents a misspecified drift function but correctly specified noise function. Suppose that the observed process satisfies

$$dX_t = -X_t dt + \sqrt{2}dW_t,$$

where  $X_0$  has a  $N(0, 1)$  distribution, the stationary distribution of  $(X_t)$ .

Estimates are calculated from the parametric model

$$dX_t = -\theta X_t^3 dt + \sqrt{2}dW_t.$$

The parameter  $\theta^*$  which minimizes  $E(X_0 - \theta X_0^3)^2$  is given by

$$\theta^* = \frac{EX_0^4}{EX_0^6} = \frac{1}{5}.$$

By Theorem 2.1,  $\hat{\theta}_T \rightarrow \frac{1}{5}$  a.s. as  $T \rightarrow \infty$ . We also have  $g(x, \theta) = \frac{1}{5}\theta^2 x^6 - 3\theta x^2$ , so that  $g'(x, \theta^*) = \frac{1}{5}x^6 - 3x^2$ . Using repeated integration by parts

$$\int_{-\infty}^s g'(z, \theta^*) d\mathbf{m}(z) = \int_{-\infty}^s \left(\frac{1}{5}z^6 - 3z^2\right) e^{-z^2/2} dz = -\left(\frac{1}{5}s^5 + s^3\right) e^{-s^2/2}$$

so that

$$\int_y^0 \int_{-\infty}^s g'(z, \theta^*) d\mathbf{m}(z) dp(s) = \frac{1}{30} y^6 + \frac{1}{4} y^4$$

and

$$\begin{aligned} \int_{-\infty}^{\infty} g'(y, \theta^*) \int_y^0 \int_{-\infty}^s g'(z, \theta^*) d\mathbf{m}(z) dp(s) dv(y) &= \int_{-\infty}^{\infty} \left(\frac{1}{5}y^6 - 3y^2\right) \left(\frac{1}{30}y^6 + \frac{1}{4}y^4\right) dv(y) \\ &= \frac{1}{150} EX_0^{12} + \frac{1}{20} EX_0^{10} - \frac{1}{10} EX_0^8 - \frac{3}{4} EX_0^6 \\ &= 94.8. \end{aligned}$$



Also

$$\int_{-\infty}^{\infty} g''(x, \theta^*) dv(x) = EX_0^6 = 15.$$

Thus  $\Sigma = \frac{(2)(94.8)}{(15)^2} = .84$ , and by Theorem 2.2 we have  $T^{1/2}(\hat{\theta}_T - \frac{1}{5}) \xrightarrow{D} N(0, .84)$ .

The asymptotic variance of  $\hat{\theta}_T$  is less than in the correctly specified case for which  $T^{1/2}(\hat{\theta}_T - 1) \xrightarrow{D} N(0, 1)$ .

2. Our second example has a correctly specified drift function and a misspecified noise function. The observed process is the same as in the first example but the parametric model is given by

$$dX_t = -\theta X_t dt + \left( \frac{2}{1+X_t^2} \right)^{1/2} dW_t.$$

Theorem 2.3 yields  $T^{1/2}(\hat{\theta}_T - 1) \xrightarrow{D} N(0, 2.75)$ . The asymptotic variance of  $\hat{\theta}_T$  has almost tripled due to the misspecified noise.

### 3. Discriminating between separate families of drift functions.

Let  $(X_t)$  satisfy (2.1) and assume throughout this section  $\sigma(x) \equiv 1$ . Suppose that two parametric models for this process have been suggested. It is required to decide in favor of the model which best fits the observed trajectory  $\{X_t, 0 \leq t \leq T\}$ .

Let  $\{f_1(x, \theta): \theta \in \Theta\}$ ,  $\{f_2(x, \phi): \phi \in \Phi\}$  be distinct families of drift functions, where  $\Theta$ ,  $\Phi$  are closed bounded intervals. A reasonable way to compare the goodness of fit of these families to the true drift function  $b(x)$  is to estimate the parameter

$$\Delta = E[f_2(X_0, \phi^*) - b(X_0)]^2 - E[f_1(X_0, \theta^*) - b(X_0)]^2.$$

In this section we introduce an estimator for  $\Delta$ . The noise function is assumed to be correctly specified; that is  $\gamma(x) \equiv 1$ .

Let  $l_T^{(1)}(\theta)$ ,  $l_T^{(2)}(\phi)$  denote the log likelihoods for the two models.

Define

$$\hat{\Delta}_T = \frac{2}{T} [l_T^{(1)}(\hat{\theta}_T) - l_T^{(2)}(\hat{\phi}_T)].$$

Given that  $f_1, f_2$  satisfy conditions (C1)-(C3), it follows from (2.7) in the proof of Theorem 2.1 that  $\hat{\Delta}_T \rightarrow \Delta$  a.s. as  $T \rightarrow \infty$ . The following result shows that  $\hat{\Delta}_T$  is asymptotically normal. Conditions from section 2 are used interchangeably between the two families of drift functions indexed by  $\theta$  and  $\phi$ .

Theorem 3.1. Suppose that  $f_1, f_2$  satisfy conditions (C1)-(C6) and (C10), where  $g_1, g_2$  are given by (2.5) with  $f = f_1, f_2$  respectively,  $\gamma(x) \equiv 1$ . Then  $T^{1/2}(\hat{\Delta}_T - \Delta) \xrightarrow{D} N(0, \Sigma_2)$ , where

$$\Sigma_2 = \frac{2}{M} \int_{-\infty}^{\infty} [g_2(y, \phi^*) - g_1(y, \theta^*) - \Delta] \int_y^0 \int_s^{\infty} [g_2(z, \phi^*) - g_1(z, \theta^*) - \Delta] dm(z) dp(s) dm(y).$$

Proof. From (2.8)

$$\begin{aligned} T^{1/2}(\hat{\Delta}_T - \Delta) &= T^{-1/2} \int_0^T [g_2(X_t, \hat{\phi}_T) - g_1(X_t, \hat{\theta}_T) - \Delta] dt \\ (3.1) \quad &+ 2T^{-1/2} [G_1(X_T, \hat{\theta}_T) - G_1(X_0, \hat{\theta}_T) - G_2(X_T, \hat{\phi}_T) + G_2(X_0, \hat{\phi}_T)]. \end{aligned}$$

The second term on the right hand side of (3.1) converges to 0 in probability.

The first term is written

$$\begin{aligned}
 & T^{-\frac{1}{2}} \int_0^T [g_2(X_t, \phi^*) - g_1(X_t, \theta^*) - \Delta] dt \\
 & + T^{-\frac{1}{2}} \int_0^T [g_2(X_t, \hat{\phi}_T) - g_2(X_t, \phi^*)] dt \\
 & + T^{-\frac{1}{2}} \int_0^T [g_1(X_t, \theta^*) - g_1(X_t, \hat{\theta}_T)] dt \\
 & = A_T + B_T + C_T.
 \end{aligned}$$

From (2.9),

$$E[g_2(X_0, \phi^*) - g_1(X_0, \theta^*) - \Delta] = 0$$

so, by Mandl (1968, p. 94),  $A_T \xrightarrow{D} N(0, \Sigma_2)$ . Next, consider  $C_T$ . Expanding  $g_1$  in a neighborhood around  $\theta^*$ ,

$$g_1(x, \theta) = g_1(x, \theta^*) + (\theta - \theta^*) g'_1(x, \bar{\theta}),$$

where  $|\bar{\theta} - \theta^*| \leq |\theta - \theta^*|$ . This gives

$$C_T = T^{\frac{1}{2}}(\hat{\theta}_T - \theta^*) \cdot \frac{1}{T} \int_0^T g'_1(X_t, \bar{\theta}_T) dt.$$

But,

$$\begin{aligned}
 \frac{1}{T} \int_0^T g'_1(X_t, \bar{\theta}_T) dt &= \frac{1}{T} \int_0^T g'_1(X_t, \theta^*) dt \\
 (3.2) \quad &+ \frac{1}{T} \int_0^T [g'_1(X_t, \bar{\theta}_T) - g'_1(X_t, \theta^*)] dt.
 \end{aligned}$$

By the proof of Theorem 2.2,  $E g'_1(X_0, \theta^*) = 0$ , so by the ergodic theorem the first term in (3.2) converges to 0 a.s. as  $T \rightarrow \infty$ . From condition (C10),

the second term in (3.2) is bounded above by

$$\phi(\bar{\theta}_T - \theta^*) \leq \frac{1}{T} \int_0^T U(X_t) dt,$$

which converges to 0 a.s. since  $|\bar{\theta}_T - \theta^*| \leq |\hat{\theta}_T - \theta^*|$ ,  $\hat{\theta}_T \rightarrow \theta^*$  a.s. and  $\frac{1}{T} \int_0^T U(X_t) dt \xrightarrow{a.s.} EU(X_0)$  as  $T \rightarrow \infty$ . Thus  $C_T \rightarrow 0$  a.s. and similarly  $B_T \rightarrow 0$  a.s..

This completes the proof of the theorem.  $\square$

### Example.

Consider the two models

$$(3.3) \quad dX_t = -\theta X_t dt + dW_t,$$

$$(3.4) \quad dX_t = -\phi X_t^3 dt + dW_t,$$

and suppose that the observed process satisfies (3.3) with  $\theta = \theta_0 > 0$ . Some involved but routine calculations give that  $\Delta = .2\theta_0$  and  $\Sigma_2 = .64\theta_0 + 9.36\theta_0^{-3}$ . Note that  $\Sigma_2 \rightarrow \infty$  as  $\theta_0 \rightarrow 0$ . The poor performance of  $\hat{\Delta}_T$  for small  $\theta_0$  is to be expected since, as  $\theta_0 \rightarrow 0$ , the drift function has less effect on the dynamics of the process so it is harder to discriminate between the two models.

### Acknowledgement.

The author thanks Loren Cobb for some helpful conversations.

# REFERENCES

- Banon, G. (1978). Nonparametric identification for diffusion processes. Siam J. on Control and Optimization 16, 380-395.
- Berger, R.L. and Langberg, N.A. (1981). Linear least squares estimates and nonlinear means. Florida State University Statistics Report M-573.
- Brown, B.M. and Hewitt, J.I. (1975). Asymptotic likelihood theory for diffusion processes. J. Appl. Prob. 12, 228-238.
- Geman, S. (1980). An application of the method of sieves: functional estimator for the drift of a diffusion. Colloquia Mathematica Societatis János Bolyai 32. Nonparametric Statistical Inference, Budapest (Hungary).
- Huber, P.J. (1967). The behavior of maximum likelihood estimates under non-standard conditions. Proc. Fifth Berkeley Symp. Math. Statist. and Prob. 1, 221-233.
- Kutoyants, Yu. A. (1977). Estimation of the drift coefficient parameter of a diffusion process in the smooth case. Theory Prob. Appl. 22, 399-406.
- Lanska, V. (1979). Minimum contrast estimation in diffusion processes. J. Appl. Prob. 16, 65-75.
- Mandl, P. (1968). Analytical Treatment of One-Dimensional Markov Processes. Academia, Prague; Springer-Verlag, Berlin.
- Prakasa Rao, B.L.S. and Rubin, H. (1981). Asymptotic theory of estimation in non-linear stochastic differential equations. Sankhyā Ser. A, to appear.
- White, H. (1981). Consequences and detection of misspecified nonlinear regression models. J. Amer. Statist. Assoc. 76, 419-433.
- Liptser, R.S. and Shirayev, A.N. (1977). Statistics of Random Processes I. Springer-Verlag, New York.

SECURITY CLASSIFICATION OF THIS PAGE

REPORT DOCUMENTATION PAGE

1. REPORT NUMBER FSU M653 USARO D-59	2. GOVT ACCESSION NO. <b>A128 799</b>	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and subtitle)		5. TYPE OF REPORT & PERIOD COVERED Technical
6. AUTHOR(s) Ian W. McKeague		6. PERFORMING ORG. REPORT NUMBER FSU Statistics Report No. M653
7. PERFORMING ORGANIZATION NAME AND ADDRESS Department of Statistics Florida State University Tallahassee, FL 32306		8. CONTRACT OR GRANT NUMBER(s) USARO Grant DAAG29-82-K-0168
9. CONTROLLING OFFICE NAME AND ADDRESS U.S. Army Research Office - Durham P.O. Box 12211 Research Triangle Park, NC 27709		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS
11. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		12. REPORT DATE March, 1983
		13. NUMBER OF PAGES 17
		15. SECURITY CLASS. (of this report) Unclassified
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE

1. DISTRIBUTION STATEMENT (of this report)

Distribution unlimited.

1. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from report)

1. SUPPLEMENTARY NOTES

THE VIEW, OPINIONS, AND/OR FINDINGS CONTAINED IN THIS REPORT ARE THOSE OF THE AUTHOR(S) AND SHOULD NOT BE CONSTRUED AS AN OFFICIAL DEPARTMENT OF THE ARMY POSITION, POLICY, OR DECISION, UNLESS SO DESIGNATED BY OTHER DOCUMENTATION.

1. KEY WORDS

Diffusion processes, maximum likelihood estimates, misspecified models.

1. ABSTRACT (Continue on reverse side if necessary and identify by block number)

The asymptotic behavior of the maximum likelihood estimator of a parameter in the drift term of a stationary ergodic diffusion process is studied under conditions in which the true drift function and true noise function do not coincide with those specified by the parametric model.

ME  
8